The following are 911 field Standing Orders (SO) for OCEMS Accredited paramedic personnel:

Based on paramedic judgment, the following may be initiated at anytime:

1. *Base Hospital contact.*
2. *Use any BLS Standing Order.*
3. *Cardiac monitor.*
4. *IV or saline lock vascular access.*

   For patients with lungs clear to auscultation and suspected poor perfusion due to hemorrhage, nausea/vomitng/diarrhea, or dehydration:

   ► *Adult/Adolescent: Normal saline bolus 250 mL; may repeat twice to maintain perfusion.*

   ► *Pediatric: 20 mL/kg (maximum 250 mL); may repeat twice to maintain perfusion.*

5. *IO for extremist patients for whom IV access cannot be established.*
6. *12-lead electrocardiogram.*
7. *Oxygen by mask or nasal canula.*
8. *BVM assisted ventilation.*
9. *Endotracheal intubation with confirmation of proper placement.*
10. *Pulse oxymetry[1]; if oxygen saturation less than 95%, give:*

    ► *High-flow oxygen by mask or nasal canula as tolerated.*

    [1]Administer oxygen by mask or nasal canula for potentially hypoxic patients when pulse oximetry may be inaccurate (hypotension, hypovolemia, hypothermia, nail polish or artificial nails, nail disease, or suspected pulse oximetry malfunction)

    ► *If patient already on home or portable oxygen, maintain flow rate and delivery (nasal canula or mask) at the rate patient is already using.*

11. *Obtain blood glucose determination and if less than 80, administer one of:*

    Adult/Adolescent:

    ► *Oral glucose preparation, if airway reflexes are intact.*

    ► *50% Dextrose 50 mL IV, may repeat once if blood glucose remains < 80.*

    ► *Glucagon I mg IM if unable to establish IV.*

    Pediatric:

    ► *Oral glucose preparation, if airway reflexes are intact.*

    ► *25% Dextrose 2 mL/kg IV if less than 2 years-old, repeat once if blood glucose remains < 80.*

    ► *50% Dextrose 1 mL/kg IV if greater than or equal to 2 years-old, repeat once if blood glucose remains < 80.*

OCEMS copyright © 2013

► *Glucagon 0.5 mg IM if unable to establish IV.*

12. For on-going seizure activity:

► Adult/Adolescent: Midazolam 5 mg IN/IM; after 3 minutes, may repeat 5 mg IN/IM once for continued or recurrent seizure activity.
► Pediatric: 0.1mg/kg IN/IM; after 3 minutes, may repeat 0.1 mg/kg IN/IM once for continued or recurrent seizure activity.

OCEMS copyright © 2013